Original Research Article

DOI: http://dx.doi.org/10.18203/2349-3291.ijcp20182577

Food allergy and allergens associated with bronchial asthma among school children in an urban and rural area of Puducherry, India: a cross-sectional study

T. R. Gobinaath, L. Anand Arokiaraj*

Department of Paediatrics, Sri Venketeswaraa Medical College, Hospital and Research Center, Puducherry, India

Received: 09 May 2018 Accepted: 31 May 2018

*Correspondence: Dr. L. Anand Arokiaraj, E-mail: drdanfi@vahoo.co.ir

E-mail: drdanfi@yahoo.co.in

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Food allergy is mostly IgE-mediated which is estimated to affect 6% to 8% children and 3% to 4% adults the clinical and social impact of which takes a hard toll in early childhood. The objective is to study the magnitude of food allergy and differences in food allergens among the urban and rural school-going children.

Methods: A cross-sectional study was conducted among 350 school going children, aged 5-10 years, attending to two private schools (n=192) and two government schools (n=158) in the urban and rural field practice areas respectively and data regarding food allergens was collected using a semi-structured proforma.

Results: There was a significant higher (p<0.001) proportion of wheeze symptomatics among the urban (n=70, 44.3%) than the rural students (n=37, 19.3%). The students had higher allergies to Ice-cream (14.57%, n=51), prawn (11.14%, n=39), peanut (14.57%, n=51), garlic (14.57%, n=51), fish (11.14%, n=39) and milk (11.14%, n=39).

Conclusions: There was a higher proportion of food allergy among the urban school students compared to the rural students. The food allergens also varied significantly in their influence on food allergy among the urban and rural areas.

Keywords: Bronchial asthma, Food allergens, School children, Urban and rural

INTRODUCTION

Allergy is an abnormal response of the immune system to a foreign substance. Neither all food substances act as foreigners to human body nor all human bodies are defensive hosts to allergens. Food allergy is mostly IgE-mediated which is estimated to affect 6% to 8% children and 3% to 4% adults the clinical and social impact of which takes a hard toll in early childhood. ¹⁻⁴ Food allergy commonly presents as gastrointestinal disturbances or skin manifestations like acute utricaria, atopic dermatitis and sometimes life-threatening anaphylaxis. However, the role of food as a trigger factor for asthma remains

vastly unexplored. Food-induced symptoms occur in approximately 2% to 29% of children and about less than 1% of adults with asthma.⁵ Food sensitization in early infancy could lead to the development of respiratory allergy and is a significant risk factor for asthma in 10% to 53% of cases.⁶⁻⁸ Allergic rhinitis has also become a frequent respiratory manifestation affecting 20% of food allergic population.⁸⁻¹⁰ Thus, food allergy stands as a patron in the pathogenesis of any atopy, especially asthma, the linkages of which are not understood clearly. The epidemiology of food allergy is influenced by genetic, cultural and geographical dietary influences. Severe and fatal reactions can occur at any age but those at greatest risk are adolescents and young adults with

asthma. Mapping the food allergens associated with asthma would help in preventing asthma in this age group. This study was designed to get an idea of possible food allergens and prevalence of food allergy among the young children and adults to further explore the association of those potential food allergens with asthma.

METHODS

This cross-sectional study was conducted among 350 school going children, aged 5-10 years, attending to two private schools (n=192) and two government schools (n=158) in the urban and rural field practice areas respectively. Initially an ethics committee approval and permission from the school heads was taken prior to conduct of the study from December to February 2017. All the schools in the urban and rural areas were listed and 4 schools were selected randomly. A prior intimation to the parents was passed on using the school information communication system and they were asked to accompany the students on the day of survey. Informed consent was taken from the parents. A child with symptom of wheezing or whistling in chest in the past 12 months was taken as criteria for defining asthma. A semistructured proforma containing socio-demographic information regarding the age, gender, occupation of parents, socio-economic status and history regarding food allergens and frequency of consumption was used to collect the data.

Statistical analysis

The collected data was entered and analyzed by using SPSS (Statistical Package for Social Sciences) version 20.0 for windows. Data is presented as frequencies and percentages. The association of food allergen with bronchial asthma and their significance was estimated using chi-square test. A p-value<0.05 was considered as statistically significant.

RESULTS

The study included 350 students attending to two private schools (n=192) and two government schools (n=158) in the field practice areas. Among the students (n=350), majority belonged to the 6-8 years age group (n=259, 74%). There was no significant difference in gender distribution as male students (n=183, 52.3%) were only a few numbers higher than the females (n=167, 47.7%). Majority (51.1%) of students were first children in their family. There was no family history of asthma in majority (92%) of students (Table 1).

The parents were interviewed about their child's unusual food consumption on the day of recent wheeze episode and ten common food items mentioned by them were listed by qualitative work-up and frequency estimated quantitatively. There was significant difference of food allergens between rural and urban schools (Table 2).

Table 1: Socio-demographic characteristics of the study population (n=350).

Participant characteristics	Urban n=158 (45.1)	Rural n=192 (54.9)	Total n=350 (100)	
Age (years)				
5-6 years	88 (55.7)	97 (50.5)	185 (52.9)	
7-8 years	42 (26.6)	87 (45.3)	129 (36.9)	
9-10 years	28 (17.7)	8 (4.2)	36 (10.3)	
Gender				
Male	84 (53.2)	99 (51.6)	183 (52.3)	
Female	74 (46.8)	93 (48.4)	167 (47.7)	
Birth order				
1	83 (52.5)	96 (50)	179 (51.1)	
2	49 (31)	79 (41.1)	128 (36.6)	
3	20 (12.7)	23 (12)	43 (12.3)	
Family History of asthma				
Present	15 (9.5)	13 (6.8)	28 (8)	
Absent	143 (90.5)	179 (93.2)	322 (92)	

Table 2: Food allergens associated with bronchial asthma.

	Urban n=158 (45.1)	Rural n=192 (54.9)	p- value	
Frequency of wheeze				
1-2 times	31	20	0.75	
3-4 times	28	11		
5-9 times	7	4		
>10 times	4	2		
Total	70	37	< 0.001	
Common food allergens				
Ice-cream	31 (19.6)	20 (10.4)	0.015	
Prawn	28 (17.7)	11 (5.7)	< 0.001	
Peanut	31 (19.6)	20 (10.4)	0.015	
Garlic	31 (19.6)	20 (10.4)	0.015	
Fish	28 (17.7)	11 (5.7)	< 0.001	
Milk	28 (17.7)	10 (5.2)	< 0.001	
Chocolate	14 (8.9)	20 (10.4)	0.62	
Curd	7 (4.4)	4 (2.1)	0.21	
Citrus fruits	7 (4.4)	4 (2.1)	0.21	
Banana	4 (2.5)	2 (1.0)	0.04	

Figures in () indicate percentages

DISCUSSION

In the present study food allergy was present in 14.57% (n=51) of the school children. The major symptom which was manifested was wheeze and itching when consumed the food known to cause the allergy. Kumar et al in their study reported that the history of food allergy was present 58.9% of the screened individuals which was higher than the present study. Rice (6.2%) followed by black gram (5.9%), lentil (5.5%), citrus fruits (5.3%), pea (3.8%), maize (3.8%) and banana (3.6%) were common food

allergens. In the present study, the students had higher allergies to Ice-cream (14.57%, n=51), prawn (11.14%, n=39), peanut (14.57%, n=51), garlic (14.57%, n=51), fish (11.14%, n=39) and milk (11.14%, n=39). The previous study used skin prick-test (SPT) to identify the food allergens whereas present study used interview method which might be the reason for difference in food allergens from the former. In another study conducted by Wang et al, they reported that sensitivity to soy, wheat, peanut, fish and egg was significantly correlated with sensitisation to some aeroallergens.⁹ The urban-rural comparison done in present study revealed that Icecream, peanut, garlic, fish, prawn were major urban food allergens whereas in rural schools Chocolates, peanuts, banana, fish were reported as allergens. There was a significant difference (p<0.001) in the proportion of wheeze symptomatics between the urban (n=70, 44.3%) and rural areas (n=37, 19.3%). The higher incidence of allergy in urban compared to rural areas may be confounded by the effect of pollution in the environment which reconfirmed the findings of Wang et al documenting that food sensitisation strongly correlated with aeroallergen sensitisation.9 Sharma et al reported egg whole, black gram and lemon as common food allergens in an adult peri-urban population which were common foods.¹² Thus, the urban allergen profile is subject to environmental confounding which needs further exploration.

CONCLUSION

The study estimated that there was a higher proportion of food allergy among the urban school students compared to the rural students. The food allergens also varied significantly in their influence on food allergy among the urban and rural areas. Further exploratory skin prick test oriented large-scale studies are needed to identify the food allergens and their pattern of affecting a particular geographical area.

Funding: No funding sources Conflict of interest: None declared

Ethical approval: The study was approved by the

Institutional Ethics Committee

REFERENCES

- 1. Sampson HA. Update on food allergy. J Allergy Clin Immunol. 2004;113:805-19.
- Osterballe M, Hansen TK, Mortz CG, Host A, Bindslev-Jensen C. The prevalence of food

- hypersensitivity in an unselected population of children and adults. Pediatr Allergy Immunol. 2005;16:567-73.
- 3. Zuberbier T, Edenharter G, Worm M, Ehlers I, Reimann S, Hantke T, et al. Prevalence of adverse reactions to food in Germany: a population study. Allergy. 2004;59:338-45.
- 4. Roehr CC, Edenharter G, Reimann S, Ehlers I, Worm M, Zuberbier T, et al. Food allergy and non-allergic food hypersensitivity in children and adolescents. Clin Exp Allergy. 2004;34:1534-41.
- 5. James JM. Respiratory manifestations of food allergy. Pediatr. 2003;111:1625-30.
- 6. Tariq SM, Matthews SM, Hakim EA, Arshad SH. Egg allergy in infancy predicts respiratory allergic disease by 4 years of age. Pediatr Allergy Immunol. 2000;11:162-7.
- Roberts G, Patel N, Levi-Schaffer F, Habibi P, Lack G. Food allergy as a risk factor for lifethreatening asthma in childhood: a case-controlled study. J Allergy Clin Immunol. 2003;112:168-74.
- 8. Penard-Morand C, Raherison C, Kopferschmitt C, Caillaud D, Lavaud F, Charpin D, et al. Prevalence of food allergy and its relationship to asthma and allergic rhinitis in schoolchildren. Allergy. 2005;60:1165-71.
- 9. Wang J, Visness CM, Sampson HA. Food allergen sensitization in inner-city children with asthma. J Allergy Clin Immunol. 2005;115:1076-80.
- 10. Bozkurt B, Karakaya G, Kalyoncu AF. Food hypersensitivity in patients with seasonal rhinitis in Ankara. Allergol Immunopathol. 2005;33:86-92.
- Kumar R, Kumari D, Srivastava P, Khare V, Fakhr H, Arora N, et al. Identification of IgE-mediated food allergy and allergens in older children and adults with asthma and allergic rhinitis. 2010. Available at: http://imsear.li.mahidol.ac.th/handle/123456789/138 636
- 12. Sharma P, Gupta P. Association of bronchial asthma and allergic rhinitis with IgE mediated allergy to common food allergens: Int J Med Res Rev. 2016;4(4):650-4.

Cite this article as: Gobinaath TR, Arokiaraj LA. Food allergy and allergens associated with bronchial asthma among school children in an urban and rural area of Puducherry, India: a cross-sectional study. Int J Contemp Pediatr 2018;5:1623-5.