

## Original Research Article

# Verbal autopsy to determine the timing and causes of infant deaths in the northern state of India

Anupam Parashar, Deepesh Barall\*, Anmol Gupta, Tripti Chauhan, Devender Kumar, Arvind Kumar, Sanjay Kumar, Gopal Ashish Sharma, Sunita Kumari

Department of Community Medicine, Indira Gandhi Medical College Shimla, Himachal Pradesh University, Himachal Pradesh, India

**Received:** 04 January 2017

**Accepted:** 30 January 2017

**\*Correspondence:**

Dr. Deepesh Barall,

E-mail: [deepeshradha2728@gmail.com](mailto:deepeshradha2728@gmail.com)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** Data on timing and probable causes of infant deaths are very important for the design of intervention to reduce mortality because of very limited study in the state. This study was undertaken to determine the timing and causes of infant deaths in our state using verbal autopsy. The suggestions and recommendations can be used to strengthen and improve the existing facilities and services for the better outcome of life in the early childhood.

**Methods:** A cross sectional study was conducted in 180 selected clusters by WHO cluster sampling method, using validated verbal autopsy tool for timing and causes of infant deaths.

**Results:** Verbal autopsy was done for 45 infant deaths. 37.8% infant deaths were early neonatal, 15.5% in late neonatal and 46.7% in post neonatal period with mean age of  $5.79 \pm 6.35$  days in neonatal period while it was  $5.07 \pm 3.38$  months in post neonatal period. Of all the infants 57.78% were females. The major causes of death were severe infection (15.5%) in neonates, pneumonia (13.3%) in post neonates followed by LBW with prematurity (11.1%) and diarrhea (11.1%). No specific cause could be determined in 13.3% cases.

**Conclusions:** The major causes of infant deaths are preventable. It emphasizes for the robust reporting mechanism to determine the exact causes of all infant mortality, strengthening of comprehensive health care delivery and ensuring the accessibility to the health care services to have a healthy infancy.

**Keywords:** Infant mortality, Neonatal mortality, Verbal autopsy

### INTRODUCTION

The Infant mortality rate of the nation is a widely accepted and long standing indicator of well being of its people. It measures the probability of dying before the age of one and the rate is expressed per 1000 live births. A high IMR is indicative of unmet health needs and unfavorable environmental factors.<sup>1</sup> The pattern of infant deaths also assesses the impact of various intervention programs aimed at improving child survival. Every year, there are an estimated 4.5 million infant deaths globally, accounting for almost 75% of deaths in children younger

than 5 years.<sup>2</sup> About a quarter of global infant deaths occur in India, which has an infant mortality of 43 per thousand live births, and it is 32 per 1000 for Himachal Pradesh, India.<sup>3</sup>

The infant mortality has almost halved worldwide in the last 25 years with India realizing impressive gains in child survival over the last two decades. However, India couldn't meet the targets for Millennium Development Goal (MDG-4), which aimed to reduce the under-five mortality rate by two thirds, between 1990 and 2015.<sup>4</sup> Also there is a small or no decline in the rate of early

neonatal mortality (ENMR), which is an indicator of quality of perinatal care.<sup>5</sup> Therefore, it is crucial to address the timing and causes of infant mortality to reduce it in India and Himachal Pradesh.

In India and other South East Asian countries, the main causes of infant mortality are thought to be preterm birth, sepsis or pneumonia, birth asphyxia, congenital abnormalities and diarrhea.

Verbal autopsies, which are interviews with care givers, have been used to establish causes of infant deaths. Data on timing and probable causes of infant deaths are very important for the design of intervention to reduce mortality as no study of infant mortality has been reported from the state.

Hence we proposed to determine the timing and causes of infant deaths in our state using verbal autopsy. The outcome of this study has been used to give suggestions and recommendations so that the existing facilities and services can be strengthened and improved for the better outcome of life in the early childhood. It will ultimately add to the overall socioeconomic development of our state and country.

## METHODS

A cross sectional study was conducted in the six districts of Himachal Pradesh for determination of timing and causes of infant deaths using verbal autopsy by the department of community Medicine, Indira Gandhi Medical College, Shimla under the aegis of National Health Mission of the state.

The period of the study was from July 2015 to September 2015.

The study was undertaken as an adjunct to the immunization survey of Himachal Pradesh using the World Health Organization's 30 X 7 cluster sampling method, developed in 1978, for the selection of clusters. A total of six districts were selected for the study with equal representation given to three zones based on altitude. Lower zone districts were Bilaspur and Mandi (altitude: 250-1000 metres), Middle zone districts were Solan and Sirmour (altitude less than 2000 metres) and upper zone districts were Shimla and Kinnaur (altitude above 2000 metres). Same clusters assigned for immunization survey were observed for the determination of timing and cause of infant deaths using the verbal autopsy. Within the cluster, all the infant deaths noticed during the preceding one year from 01 July 2014 to 30 June 2015 were included 6.

### *Inclusion criteria*

- All the infants who died during the specified period and the parents or family members were available and gave consent.

- Infant belonged to the selected cluster.

From each of the 6 districts, 30 clusters using the WHO 30 X 7 cluster sampling method were selected. In each cluster, enquiry was done from Health worker/AWW/ASHA worker or any responsible village personnel for any infant deaths during the specified period in that cluster area. The household was visited by the team of Medical Officers and after confirming the eligibility criteria and consent from the concerned family member, verbal autopsy form 9, as used in the million-death study by Registrar General of India through Centre of Global Health Research 10 was filled with the information as required along with the detailed narration of events that led to death, from the relative (preferably mother) of the deceased infant. A detailed history of the events of birth and the circumstances leading to death were elicited from the respondent. The age of the deceased child was ascertained by the exact date of birth if the parents could recall or by the religious and the ritual events. Subsequent cases were searched using snowball sampling technique or if already in the notice of the visited family or health worker /ASHA/ Aanganwadi worker of that area. The stillbirths were excluded from the study. The cause of death was ascertained using standard verbal autopsy procedure. In case of doubt, the cause of death was determined after discussion with the jury of consultants in the Department of Community Medicine at IGMC Shimla.

The study was approved by the Institutional Ethics Committee of IGMC Shimla.

## RESULTS

In this study, six teams visited 21146 households in 180 clusters, where we came across 45 infant deaths. We excluded the four still births that came across in our study.

Interview was taken from mothers of the deceased child in 57.78% cases; from grandparents in 22.23% and 20% were other relatives.

37.78% infant deaths were early neonatal out of which 20% were on the first day of life (maximum infant deaths), 15.5% in late neonatal and 46.67% in post neonatal period. Mean age of death in neonatal period was 5.79 days while in post neonatal period, it was 5.07 months. Half of the deaths in the neonatal period occurred by the end of 3rd day while half of the deaths in the post neonatal period occurred by the end of 5th month.

Overall there were 57.78% female and 42.23% male among infant deaths whereas the neonatal mortality was found equal for both the sexes in our study.

We observed the cause of death in infants to be in the following order of magnitude: severe infection (15.55%) in neonates, pneumonia (13.33%) in post neonates followed by LBW with prematurity (11.11%) and diarrhea (11.11%), then by birth asphyxia (8.89%) and congenital anomalies (8.89%) and then injuries (2.22%). In 15.55% cases, there were other causes and in 13.33% cases, cause couldn't be determined. Duration of illness before death ranged from few hours in neonates to 25 days in post neonate.

Other significant findings of our study include 62.5% of neonates were delivered at some health institution while 33.3% at home conducted by either trained or untrained TBA. 42.23% of children died at some health institutions and 40% at home. 65% babies were of normal size and weight at time of birth while it was low in 16% with multiple pregnancies as a cause of preterm and LBW in

12% of cases and high in 12 % of cases. Almost 85% post neonates were vaccinated for their age.

## DISCUSSION

In this study, we observed 45 infant deaths out of which 53% deaths were in the neonatal period (37.78% in the early neonatal period and 15.5% in the late neonatal period) as compared to 60-68.5% in previous studies.<sup>11,12</sup> It is also below our national and state figure of 69% and 77.5% respectively as per SRS 2012. The lower proportion of neonatal deaths in our study may be true, supported by the adequate antenatal health coverage reflecting the maternal health status during the pregnancy and better socioeconomic conditions reflected by per capita income in our state as compared to the rest of the country (Table 1).

**Table 1: District wise demographic and facility details (SRS 2014).**

	Shimla	Sirmour	Bilaspur	Kinnaur	Solan	Mandi
Total population (in thousands)	814	529	382	84.1	580.3	999.8
Average literacy rate (%age)	83.64	78.80	84.59	80	83.68	81.53
Female literacy rate (%age)	77.13	71.36	77.9	70.96	76.97	73.66
Total clusters covered	30	30	30	30	30	30
Total household covered (Numbers)	4203	2986	2805	4415	3352	3385
Total infant deaths encountered (Numbers)	5	7	7	9	7	10
Villages having Govt. health facility(%age) <sup>7</sup>	70	69.4	75.7	72.5	60.6	48.6
Per capita income of Himachal (Rs.)	130,067 (2015-16) <sup>8</sup>					

The higher proportion of 46.67% deaths in the post neonatal period as compared to national (31%), state (22.5%) and other study (40%) can be due to the casual health seeking behavior of the people for perceived presence of danger signs despite available preventive and treatment facilities.<sup>13,14</sup> Or the difference can be merely due to small sample size of our study.

In this study the gender distribution of deaths in infancy was 57.78% in females and 42.23% in males which is almost similar to what had been observed in previous study (53 and 47% respectively).<sup>15</sup>

In this study the main causes of death in neonates were severe infection (29%) followed by LBW with prematurity (21%), Birth Asphyxia(16.6%), Congenital anomalies (12.5%) and diarrhea (4%). In post neonates, deaths were due to Pneumonia in (28.5%) followed by Diarrhoea 19% (4), Injury 4.7% (1) and congenital anomalies 4.7%(1). These findings were coherent with results from the community based studies in South Asia which have reported neonatal causes of death within the following ranges: preterm birth, 8-38%; sepsis or pneumonia, 7-52%; birth asphyxia 10-28%; (sometimes

including birth injury); tetanus 2-36%; and diarrhea 9%.<sup>16-18</sup>

We were unable to find cause for 13.34% of deaths which is within the range of previous study who has reported similar proportions of death for which no specific cause could be identified by verbal autopsy.<sup>19</sup>

In this study 42.23% of deaths were in the health institution while 40% were at home. So in settings where considerable number of deaths occurs outside of the health system, verbal autopsy is considered the most practicable method for determining the cause of death but with all its limitation as a part of study. These includes problems inherent to certain case definitions, difficulty in eliciting closely related symptoms as in severe pneumonia and septicemia, limited in differentiating viral and bacterial infections, subjectively classifying preterm babies of their sizes and weight and limited information regarding the maternal health status and the last treatment offered to the deceased child. Although every effort has been done to overcome all the limitations but some unforeseen circumstances are always unavoidable.

The strength of this study is that it used the previously validated methods for both sampling as well as for gathering information i.e. standardized and well validated verbal autopsy tool as used in the million-death study by the Registrar General of India through Centre of Global Health Research. Further the whole study was carried by the team of doctors, all of whom are the postgraduates in the community medicine department having exposure and experience in the related subject and above that the causes of death were further confirmed by the independent team of consultants in the department of community medicine.

The process of examining deaths by number of days since birth yielded useful insights for health program planning and illustrates the need for a continuum of care. Antenatal education regarding health and nutritional status of the mother, routine antenatal visits, clean delivery practices, immediate breast feeding and care seeking for complications is required. In view of high deaths on the first day of life, it emphasizes the need to increase the coverage of skilled birth attendants and to equip them with the capabilities to identify and manage birth complications including birth asphyxia. For early deaths in the post-natal period, it also requires to ensure strict adherence for vigilant postnatal visits with preventive care for all newborn babies and prompt recognition and referral for the needy along with universal access to quality emergency neonatal care.

Study findings suggest that the surveillance during the pregnancy and postnatal period needs to be effectively implemented and monitored. A key finding of this study is that the burden of early neonatal death is high for which more detailed and extensive studies are required to define the exact burden and to evaluate the ongoing programs on continuum of care. And greater coverage of antepartum, intrapartum and early postnatal health interventions, in combination with promotion of care seeking behavior and effective communication between community and health facilities would be expected to have the greatest effect in this setting.

## CONCLUSION

This study which is one of the few of its kind in the state resulted in the determination of the timing and causes of infant mortality for its comparison to other parts of the country. The major causes of infant deaths are preventable to a large extent. Although our state has improved statistically in the indicators of economic growth, literacy and infant mortality over a period of time but a lot of improvement is yet to be done at root level, especially in the determination of exact causes of all infant mortality with robust reporting mechanism and delivery of comprehensive health care services to all newborns and infants. The burden of mortality among infants can be reduced by strengthening of continuum of care through RMNCH+A program, mainly promotion of institutional deliveries, intranatal and neonatal care, early

recognition of danger signs at household and facility level and correct management through trained health functionaries. Also it emphasizes the need to ensure the accessibility to the health care services along with strengthening of referral system to have a healthy infancy to all born alive.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Institutional Ethics Committee*

## REFERENCES

1. NIMS, ICMR and UNICEF, infant and child Mortality in India: Levels, Trends and Determinants, National Institute of Medical Statistics (NIMS), Indian Council of Medical Research(ICMR), and UNICEF India Country Office, New Delhi, India, 2012.
2. Global health observatory (GHO) data. Available at [http://www.who.int/gho/child\\_health/mortality/neonatal\\_infant\\_text/en/](http://www.who.int/gho/child_health/mortality/neonatal_infant_text/en/). Accessed on 02 September 2016.
3. Registrar General India. Sample Registration System. SRS bulletin, September 2014 released on July 2016. Available at [http://censusindia.gov.in/vital\\_statistics/SRS\\_Bulletins/SRS%20Bulletin%20-September%202014.pdf](http://censusindia.gov.in/vital_statistics/SRS_Bulletins/SRS%20Bulletin%20-September%202014.pdf) Accessed on 02 September 2016.
4. The Infant and Child Mortality India Report Available at <http://unicef.in/PressReleases/374/The-Infant-and-Child-Mortality-India-Report#sthash.eiHwz6QT.dpuf>. Accessed on 02 September 2016.
5. National Institute of Medical Statistics (NIMS), Indian Council of Medical Research (ICMR), and UNICEF India Country Office .Infant and Child Mortality in India, Levels Trends and determinants, Fact sheet, New Delhi, India; 2012.
6. Government of Himachal Pradesh, India. Available at <http://himachal.nic.in/>. Accessed on 02 September 2016.
7. International Institute for Population Sciences (IIPS). District Level Household and Facility Survey (DLHS-4), 2012-13: India. Himachal Pradesh: Mumbai: IIPS; 2014.
8. Economic survey projects Himachal Pradesh's growth rate at 7.7% - Times of India". Available at <http://timesofindia.indiatimes.com/city/shimla/Economic-Survey-projects-Himachal-Pradeshs-growth-rate-at-7-7/articleshow/51266064.cms>. Accessed on 02 September 2016.
9. Anker M, Black R, Coldham C, Kalter HD, Quigley MA, Ross D, et al. A standard verbal autopsy method for investigating causes of death in infants and children. Geneva, World Health Organization, 1999. Available at [http://whqlibdoc.who.int/hq/1999/WHO\\_CDS\\_CSR\\_ISR\\_99.4.pdf](http://whqlibdoc.who.int/hq/1999/WHO_CDS_CSR_ISR_99.4.pdf). Accessed on 02 September 2016.

10. Registrar General of India, Centre of Global Health Research. Prospective Study of One Million Deaths in India: Rationale, design and validation results. Final version; 2005.
11. Rasaily R. Age profile of neonatal death short communication Indian Council of Medical Research, Ansari Nagar, New Delhi 110029, India. *Indian Pediatr.* 2008;45:991-4.
12. Shah MS, Khaliq N, Khan Z, Amir A. Verbal autopsy to determine causes of deaths among under-five children. *Curr Pediat Res.* 2010;14(1):51-3.
13. SRS bulletin 2012, Sample Registration System, Registrar General India, Available at [censusindia.gov.in/vital\\_statistics/SRS\\_Bulletins/SRS\\_Bulletin-September\\_2013.pdf](http://censusindia.gov.in/vital_statistics/SRS_Bulletins/SRS_Bulletin-September_2013.pdf). Accessed on 04 September 2016.
14. Shah MS, Khaliq N, Khan Z, Amir A. Verbal Autopsy to determine causes of deaths among under-five children. *Curr Pediat Res.* 2010;14(1):51-3.
15. Khanna R, Kumar A, Vaghela JF, Sreenivas V, Puliyl JM. Community based retrospective study of sex in infant mortality in India. *BMJ.* 2003;327:126.
16. Fikree FF, Azam SI, Berendes HW. Time to focus child survival programs on the newborn: assessment of levels and causes of infant mortality in rural Pakistan. *Bull World Health Organ.* 2002;80:271-6.
17. Freeman JV, Christian P, Khatri SK, Adhikari RK, Leclercq SC, Katz J, et al. Evaluation of neonatal verbal autopsy using physician review versus algorithm-based cause of death assignment in rural Nepal. *Paediatr Perinat Epidemiol.* 2005;19:323-31.
18. Baqui AH, Paul VK, Reddy HM, Baitule SB. Why do neonates die in rural Gadchiroli, India? (Part I): primary causes of death assigned by neonatologist based on prospectively observed records. *J Perinatol.* 2005;25 Suppl 1:S29-34.
19. Aggarwal AK, Kumar P, Pandit S, Kumar R. Accuracy of WHO verbal autopsy tool in determining major causes of neonatal deaths in India. *PLOS ONE.* 2013(8);1-9.

**Cite this article as:** Parashar A, Barall D, Gupta A, Chauhan T, Kumar D, Kumar A, et al. Verbal autopsy to determine the timing and causes of infant deaths in the northern state of India. *Int J Contemp Pediatr* 2017;4:591-5.